TECHNICAL BULLETIN

Maternal Zinc Status Supports Offspring Gut Development, Immune Function and Neurological Health

Introduction

Zinc (Zn) is an essential trace mineral required for normal growth, development and immune function. During pregnancy, Zn requirements increase substantially to support fetal growth and organ development. While Zn is well known for its role in brain and immune function, emerging evidence suggests that adequate maternal Zn status may also be critical for gastrointestinal (GI) tract development and for the long-term influence on the offspring's neurological and immune health.

The GI tract is more than a digestive organ. It serves as a critical barrier between the external environment and the body, regulates nutrient absorption and houses a large portion of the immune system. Disruptions in gut development during early life may contribute to altered nutrient absorption, immune dysregulation and changes in gut-brain communication pathways.

This technical bulletin reviews the role of Zn in intestinal development, the relationship between gut health and neurological function, and the importance of maintaining adequate Zn status during pregnancy and early life.

Zinc and Gut Development

Proper development of the intestinal tract requires tightly regulated processes including cellular proliferation, differentiation and tissue remodeling (Figure 1). Many of these processes rely on Zn-dependent enzymes, transcription factors and signaling pathways. As a result, inadequate Zn availability during development may alter intestinal structure and function.

Research has demonstrated that Zn status influences several aspects of intestinal development, including villus height, villus-to-crypt ratio, epithelial cell renewal and goblet cell abundance. These structural features contribute directly to nutrient digestion, nutrient absorption and barrier integrity. Zinc deficiency has been associated with shortened villi, reduced absorptive surface area, impaired epithelial repair and altered intestinal morphology.




Figure 1. Influence of Zn on Intestinal Development

Additionally, Zn is required for normal activity of several brush border enzymes involved in digestion and nutrient utilization. Deficiencies have been shown to reduce activity of enzymes responsible for carbohydrate digestion and nutrient processing, potentially impairing GI function.

Consequences of Zinc Deficiency During Development

The effects of Zn deficiency extend beyond intestinal morphology. Zinc deficiency has been associated with increased intestinal permeability, impaired mucosal integrity, altered mucus production and inflammation within the GI tract. These changes may reduce the intestine's ability to function as an effective barrier while simultaneously compromising nutrient absorption.

In severe cases, Zn deficiency has been associated with villus atrophy, intestinal ulceration and mucosal damage. Even moderate deficiencies may impair epithelial renewal and intestinal resilience. Because the intestine serves as the primary site of Zn absorption, compromised intestinal development may further perpetuate Zn deficiency through reduced nutrient uptake.

Collectively, these findings suggest that adequate maternal Zn status plays an important role in establishing a healthy gastrointestinal tract capable of supporting nutrient absorption, immune function and the baby's long-term health.

All trademarks are the property of Zinpro Corp.

©2026 Zinpro Corp. All rights reserved.

The Gut-Brain Connection

Growing evidence supports communication between the GI tract and the central nervous system through a network of neural, immune and endocrine pathways often described as the gut-brain axis (Figure 2). Signals originating from the GI tract can influence neurological function, while the brain can influence GI physiology through interactions with the enteric nervous system and stress-response pathways.

Disruptions in gut barrier function may increase exposure of the immune system to dietary antigens, microbial metabolites and inflammatory signals. In turn, these events may alter immune signaling pathways that influence neurological development and function. Vela et al. (2015) proposes that maternal Zn deficiency may contribute to a cycle involving GI dysfunction, immune dysregulation, stress responses and altered neurological outcomes during fetal development (Figure 2).




Figure 2. Proposed relationship between maternal Zn status, gut development, and gut-brain communication. Due to the multifaceted effects of Zn on gut development and morphology, pre– and perinatal Zn deficiency might affect neonatal gut development, and adequate Zn status potentially mitigates many dysfunctions associated with neurological disorders.

The precise contribution of Zn status to neurological disorders is an active area of scientific investigation, with current evidence supporting an important role for Zn in maintaining the integrity of interconnected gut, immune and neurological systems.

Bioavailability and Factors that May Reduce Zinc Availability During Pregnancy

Maternal Zn deficiency is not always caused by inadequate Zn intake. A variety of dietary components, nutritional supplements, and commonly used medications may reduce Zn bioavailability by interfering with Zn absorption. Phytates, folic acid, calcium, iron, copper, high-fructose corn syrup (HFCS), and certain medications have all been associated with reduced Zn availability. Because fetal development depends on maternal Zn status, these interactions may affect Zn delivery to the developing offspring even when dietary Zn intake appears sufficient.

Reduced Zn availability during pregnancy may impair fetal intestinal development, alter gut barrier integrity, affect nutrient absorption, and contribute to gastrointestinal dysfunction later in life. Several of these abnormalities, including vitamin and mineral malabsorption, metallothionein dysfunction, altered Cu:Zn relationships, and gastrointestinal disturbances, have been reported in association with autism spectrum disorder and other neurological disorders.

Zinc source should be considered when developing nutritional strategies to support maternal Zn status. Zinc amino acid complexes offer advantages under conditions where dietary antagonists are present because they can utilize amino acid-associated absorption mechanisms and may be less susceptible to competitive interactions that reduce the bioavailability of conventional inorganic Zn sources. As pregnancy places increased demands on maternal Zn reserves, maintaining adequate Zn availability is an important consideration for supporting maternal and fetal health.

Implications for Nutritional Strategies

Current evidence suggests that maternal Zn status may influence normal intestinal development, mucosal integrity, nutrient absorption, and immune function in the developing offspring. By influencing the development of systems that contribute to gut-brain communication, adequate maternal Zn nutrition may help support healthy developmental outcomes.

Nutritional strategies designed to support maternal Zn status should consider both Zn intake and bioavailability. Numerous dietary and physiological factors can reduce Zn absorption; therefore, Zn source is an important consideration when developing nutritional programs intended to support maternal and fetal health.

Source: Vela et al., 2015. Neural Plasticity Volume 2015, Article ID 972791 **doi:**10.1155/2015/972791

All trademarks are the property of Zinpro Corp.

©2026 Zinpro Corp. All rights reserved.